



HOMOTOPY INERTIA GROUPS AND TANGENTIAL STRUCTURES

Ramesh Kasilingam

Statistics and Mathematics Unit

Indian Statistical Institute

Bangalore Centre, Bangalore - 560059

Karnataka, India

e-mail: rameshkasilingam.iitb@gmail.com

mathsramesh1984@gmail.com

Abstract

We show that if M and N have the same homotopy type of simply connected closed smooth m -manifolds such that the integral and mod-2 cohomologies of M vanish in odd degrees, then their homotopy inertia groups are equal. Let M^{2n} be a closed $(n-1)$ -connected $2n$ -dimensional smooth manifold. We show that, for $n=4$, the homotopy inertia group of M^{2n} is trivial and if $n=8$ and $H^n(M^{2n}; \mathbb{Z}) \cong \mathbb{Z}$, then the homotopy inertia group of M^{2n} is also trivial. We further compute the group $\mathcal{C}(M^{2n})$ of concordance classes of smoothings of M^{2n} for $n=8$. Finally, we show that if a smooth manifold N is tangentially homotopy equivalent to M^8 , then N is diffeomorphic to the connected sum of M^8 and a homotopy 8-sphere.

Received: November 10, 2016; Revised: May 8, 2017; Accepted: May 10, 2017

2010 Mathematics Subject Classification: Primary 57R55, 57R65; Secondary 55Q45, 55Q55.

Keywords and phrases: projective plane like manifolds, smooth structures, homotopy inertia groups, cohomotopy groups.

Communicated by Michael R. Kelly

1. Introduction

Let Θ_m be the group of smooth homotopy spheres defined by Kervaire and Milnor in [18]. Recall that the collection of homotopy spheres Σ which admit a diffeomorphism $M \rightarrow M\#\Sigma$ form a subgroup $I(M)$ of Θ_m , called the *inertia group* of M , where we regard the connected sum $M\#\Sigma^m$ as a smooth manifold with the same underlying topological space as M and with smooth structure differing from that of M only on an m -disc. The homotopy inertia group $I_h(M)$ of M^m is a subset of the inertia group consisting of homotopy spheres Σ for which the identity map $id : M \rightarrow M\#\Sigma^m$ is homotopic to a diffeomorphism. Similarly, the concordance inertia group of M^m , $I_c(M^m) \subseteq \Theta_m$, consists of those homotopy spheres Σ^m such that M and $M\#\Sigma^m$ are concordant. Among these groups, only the concordance inertia group $I_c(M)$ is a homotopy type invariant of M [5, 17]. Also, in [12], it was shown that the homotopy inertia group $I_h(M)$ is an h -cobordism invariant for manifolds of dimension at least eight.

The paper is organized as following. Preliminary definitions and notations are given in Section 2. In Section 3, we show that if M and N have the same homotopy type of simply connected closed smooth m -manifolds such that the integral and mod-2 cohomologies of M vanish in odd degrees, then their homotopy inertia groups are equal. Thus, the homotopy inertia group $I_h(M)$ is a homotopy type invariant for closed oriented simply connected manifolds with vanishing the integral and mod-2 cohomology in odd degrees. In fact, we show that if M^m ($m \geq 5$) is a closed simply connected smooth m -manifold such that the integral and mod-2 cohomologies of M^m vanish in odd degrees, then the natural forgetful map $F_{Con} : \mathcal{C}^{Diff}(M) \rightarrow \mathcal{S}^{Diff}(M)$ is injective, where $\mathcal{C}^{Diff}(M)$ is the group of concordance classes of smooth structures on M^m and $\mathcal{S}^{Diff}(M)$ is the set of diffeomorphism classes of homotopy-smooth structures on M^m .

Let M^{2n} be a closed $(n-1)$ -connected $2n$ -dimensional smooth manifold. In this section, we also show that, for $n=4$, the homotopy inertia group $I_h(M^{2n})$ is trivial and if $n=8$ and $H^n(M^{2n}; \mathbb{Z}) \cong \mathbb{Z}$, then the homotopy inertia group $I_h(M^{2n})$ is also trivial.

In Section 4, we compute the 0th reduced stable cohomotopy group $\tilde{\pi}^0(M^{2n})$ for $n=4, 8$ and the group $\mathcal{C}(M^{2n})$ of concordance classes of smoothings of M^{2n} for $n=8$, which will be used in Section 5.

In Section 5, with the use of surgery theoretic structure sets, we give a diffeomorphism classification of closed smooth manifolds in the tangential homotopy type of M^{2n} . Recall that a homotopy equivalence of smooth manifolds of the same dimension $f: L \rightarrow N$ is called *tangential* if the vector bundles $f^*(T(N))$ and $T(L)$ are stably isomorphic. In [27], it was studied that under what conditions tangentially homotopy equivalent manifolds are necessarily homeomorphic. Examples of non-homeomorphic tangentially homotopy equivalent manifolds were given in [8, 10]. In this section, we show that if $n \equiv 0 \pmod{4}$, $n > 0$ and a smooth manifold N is tangentially homotopy equivalent to M^{2n} , then N is homeomorphic to M^{2n} . We also show that if a smooth manifold N is tangentially homotopy equivalent to M^8 , then N is diffeomorphic to $M^8 \# \Sigma$, where $\Sigma \in \Theta_8$ is a homotopy 8-sphere.

Notation. We denote by $[X; Y]$ the set of all free homotopy classes of maps from X to Y . If X, Y are well pointed spaces, and if Y is 0-connected, then the fundamental group $\pi_1(Y)$ acts on the set $[X; Y]_0$ of based homotopy classes; the set $[X; Y]$ of all free homotopy classes can be identified with the orbit set of this action [40, Section III.1]. If Y is an H -space (or if Y is 1-connected), this action is trivial, so $[X; Y] = [X; Y]_0$. Denote by $O = \text{colim}_n O(n)$, $PL = \text{colim}_n PL(n)$, $Top = \text{colim}_n Top(n)$, $G =$

$\text{colim}_n G(n)$, the direct limit of the groups of orthogonal transformations, PL homeomorphisms, homeomorphisms and homotopy equivalences, respectively. We also denote $SPL = \text{colim}_n SPL(n)$ and $SG = \text{colim}_n SG(n)$, the direct limit of the groups of orientation-preserving PL homeomorphisms and homotopy equivalences, respectively.

In this paper, all the manifolds will be closed, smooth, oriented and connected, and all the homotopy equivalences, homeomorphisms and diffeomorphisms are assumed to preserve orientation, unless otherwise stated.

2. Preliminaries

Definition 2.1. (a) A homotopy m -sphere Σ^m is an oriented smooth closed manifold homotopy equivalent to the standard unit sphere \mathbb{S}^m in \mathbb{R}^{m+1} .

(b) A homotopy m -sphere Σ^m is said to be *exotic* if it is not diffeomorphic to \mathbb{S}^m .

(c) Two homotopy m -spheres Σ_1^m and Σ_2^m are said to be *equivalent* if there exists an orientation preserving diffeomorphism $f : \Sigma_1^m \rightarrow \Sigma_2^m$.

The set of equivalence classes of homotopy m -spheres is denoted by Θ_m .

Definition 2.2. ($Cat = Diff$ or PL). Let M be a closed Cat -manifold. Let (N, f) be a pair consisting of a closed Cat -manifold N together with a homeomorphism $f : N \rightarrow M$. Two such pairs (N_1, f_1) and (N_2, f_2) are concordant provided there exists a Cat -isomorphism $g : N_1 \rightarrow N_2$ such that the composition $f_2 \circ g$ is topologically concordant to f_1 , i.e., there exists a homeomorphism $F : N_1 \times [0, 1] \rightarrow M \times [0, 1]$ such that $F|_{N_1 \times 0} = f_1$ and

$F_{|N_1 \times 1} = f_2 \circ g$. The set of all such concordance classes is denoted by $\mathcal{C}^{Cat}(M)$.

We will denote the class in $\mathcal{C}^{Cat}(M)$ of (N, f) by $[N, f]$. The base point of $\mathcal{C}^{Cat}(M)$ is the equivalence class $[M, Id]$ of $Id : M \rightarrow M$.

Definition 2.3. (*Cat = Diff or Top-structure sets*). Let M be a closed *Cat*-manifold. We define the *Cat*-structure set $\mathcal{S}^{Cat}(M)$ to be the set of equivalence classes of pairs (N, f) , where N is a closed *Cat*-manifold and $f : N \rightarrow M$ is a homotopy equivalence. And the equivalence relation is defined as follows:

$$(N_1, f_1) \sim (N_2, f_2) \text{ if there is a } Cat\text{-isomorphism } h : N_1 \rightarrow N_2 \\ \text{such that } f_2 \circ h \text{ is homotopic to } f_1.$$

We will denote the class in $\mathcal{S}^{Cat}(M)$ of (N, f) by $[N, f]$. The base point of $\mathcal{S}^{Cat}(M)$ is the equivalence class $[M, Id]$ of $Id : M \rightarrow M$. Denote by $[M^m \# \Sigma^m]$ the equivalence class of $(M^m \# \Sigma^m, Id)$ in $\mathcal{S}^{Cat}(M)$ or $\mathcal{C}^{Diff}(M)$. (Note that $[M^m \# S^m]$ is the class of (M^m, Id) .)

Definition 2.4. Let M^m be a closed smooth m -dimensional manifold. The inertia group $I(M) \subset \Theta_m$ is defined as the set of $\Sigma \in \Theta_m$ for which there exists a diffeomorphism $\phi : M \rightarrow M \# \Sigma$.

Define the homotopy inertia group $I_h(M)$ to be the set of all $\Sigma \in I(M)$ such that there exists a diffeomorphism $M \rightarrow M \# \Sigma$ which is homotopic to $Id : M \rightarrow M \# \Sigma$.

Define the concordance inertia group $I_c(M)$ to be the set of all $\Sigma \in I_h(M)$ such that $M \# \Sigma$ is concordant to M .

Observe that there are natural forgetful maps $F_{Con} : \mathcal{C}^{Diff}(M) \rightarrow$

$\mathcal{S}^{Diff}(M)$ and $F_{Diff} : \mathcal{S}^{Diff}(M) \rightarrow \mathcal{S}^{Top}(M)$. In [39], Weinberger showed that the forgetful map $F_{Diff} : \mathcal{S}^{Diff}(M) \rightarrow \mathcal{S}^{Top}(M)$ is finite-to-one with image containing a subgroup of finite index. It also follows that the forgetful maps $F_{Diff} : \mathcal{S}^{Diff}(M) \rightarrow \mathcal{S}^{Top}(M)$ and $F_{Con} : \mathcal{C}^{Diff}(M) \rightarrow \mathcal{S}^{Diff}(M)$ fit into a short exact sequence of pointed sets:

$$\mathcal{C}^{Diff}(M) \xrightarrow{F_{Con}} \mathcal{S}^{Diff}(M) \xrightarrow{F_{Diff}} \mathcal{S}^{Top}(M).$$

Definition 2.5 [4, 38, 24, 28, 29]. Let N and M be closed smooth manifolds. A degree one normal map $(N, f, b) : N \rightarrow M$ is a degree one map $f : N \rightarrow M$ together with a stable bundle map $b : \nu_N \rightarrow \xi$, where ν_N is the stable normal bundle of N , b covers f and ξ is some stable vector bundle over M (necessarily fibre homotopy equivalent to ν_M).

A normal bordism of degree one normal maps (N_i, f_i, b_i) , $i = 0, 1$ is a degree one normal map $(Z, g, c) : Z \rightarrow M \times [0, 1]$ restricting to (N_i, f_i, b_i) over $M \times \{i\}$.

The set of normal bordism classes of degree one normal maps into a closed smooth manifold M is called the *normal structure set* of M , and we denote it by $\mathcal{N}^{Diff}(M)$.

The normal structure set $\mathcal{N}^{Diff}(M)$ of an m -dimensional smooth closed simply connected manifold M^m fits into the surgery exact sequence of pointed sets

$$\begin{aligned} \dots \rightarrow \mathcal{N}^{Diff}(M \times [0, 1]) \xrightarrow{\sigma} L_{m+1}(\mathbb{Z}) \xrightarrow{\theta} \mathcal{S}^{Diff}(M) \\ \xrightarrow{\eta} \mathcal{N}^{Diff}(M) \xrightarrow{\sigma} L_m(\mathbb{Z}), \end{aligned} \quad (2.1)$$

where the map $\eta : \mathcal{S}^{Diff}(M) \rightarrow \mathcal{N}^{Diff}(M)$ is defined by mapping $[N, f]$ to $[(N, f, b) : N \rightarrow M]$, here $b : \nu_N \rightarrow (f^{-1})^*(\nu_M)$ is the canonical

bundle map and $\xi = (f^{-1})^*(\nu_N)$, $L_m(\mathbb{Z})$ are the surgery obstruction groups, the maps $\sigma : \mathcal{N}^{Diff}(M) \rightarrow L_m(\mathbb{Z})$ and $\sigma : \mathcal{N}^{Diff}(M \times [0, 1]) \rightarrow L_{m+1}(\mathbb{Z})$ are the surgery obstruction maps and the map $\theta : L_{m+1}(\mathbb{Z}) \rightarrow \mathcal{S}^{Diff}(M)$ is the action of $L_{m+1}(\mathbb{Z})$ on $\mathcal{S}^{Diff}(M)$. Using identity maps as base points, we have Sullivan's familiar identifications $\mathcal{N}^{Diff}(M) = [M, G/O]$ and $\mathcal{N}^{Diff}(M \times [0, 1]) = [SM, G/O]$, where SM is the suspension of M .

Definition 2.6 [27, 9]. Let M be a closed smooth m -manifold with stable normal bundle ν_M of rank $k \gg m$. The smooth tangential structure set of M :

$$\mathcal{S}^{TDiff}(M) = \{(N, f, b) \mid f : N \rightarrow M, b : \nu_N \rightarrow \nu_M\} / \simeq,$$

consists of equivalence classes of triples (N, f, b) , where N is a smooth manifold, $f : N \rightarrow M$ is a homotopy equivalence and $b : \nu_N \rightarrow \nu_M$ is a map of stable bundles. Two structures (N_0, f_0, b_0) and (N_1, f_1, b_1) are *equivalent* if there is an s -cobordism (U, N_0, N_1, F, B) and where $F : U \rightarrow M$ is a simple homotopy equivalence, $B : \nu_U \rightarrow \nu_M$ is a bundle map and these data restrict to (N_0, f_0, b_0) and (N_1, f_1, b_1) at the boundary of U .

The tangential surgery exact sequence for a simply connected closed smooth manifold M finishes with the following four terms:

$$L_{m+1}(\mathbb{Z}) \xrightarrow{\theta} \mathcal{S}^{TDiff}(M) \xrightarrow{\eta^t} \mathcal{N}^{TDiff}(M) \xrightarrow{\sigma} L_m(\mathbb{Z}), \quad (2.2)$$

where the definition of $\mathcal{N}^{TDiff}(M)$ is similar to the definition of $\mathcal{S}^{TDiff}(M)$ except that for representatives (N, f, b) we require only that $f : N \rightarrow M$ is a degree one map and the equivalence relation is defined using normal cobordisms over (M, ν_M) . The set $\mathcal{N}^{TDiff}(M)$ of tangential normal invariants of M can be identified with $[M, SG]$.

Let $QM = \Omega^\infty \Sigma^\infty M = \text{colim}_n \Omega^n \Sigma^n M$. Then QM is an infinite loop space, i.e., the zero space of the Ω -spectrum $\{Q\Sigma^n M\}$. For $M = S^0$, QS^0 has components $Q_k S^0$, $k \in \mathbb{Z}$, determined by the degree of self maps of spheres. We let $Q_k S^0$ be the k th component of QS^0 . The space QS^0 is an H -space under the loop product

$$* : QS^0 \times QS^0 \mapsto QS^0$$

which satisfies

$$* : Q_s S^0 \times Q_t S^0 \mapsto Q_{s+t} S^0,$$

and for any space M there is a free and transitive action

$$[M, Q_1 S^0] \times [M, Q_0 S^0] \xrightarrow{*} [M, Q_1 S^0]([\phi], [\alpha]) \mapsto [\phi] * [\alpha].$$

It is important to note that $Q_0 S^0$ has the same homotopy type of SG and the H -space structure on SG given by composition of maps corresponds not to the loop product on $Q_0 S^0$ [36]. Therefore, $[M, SG] = [M, Q_0 S^0] = \tilde{\pi}^0(M)$ as sets, where $\tilde{\pi}^0(M)$ is the 0th reduced stable cohomotopy group of M . Actually, $\tilde{\pi}^0(M)$ is a ring, and, as groups, $[M, SG] \cong 1 + \tilde{\pi}^0(M)$ where the addition on the right is given by $(1 + \alpha)(1 + \beta) = 1 + \alpha + \beta + \alpha\beta$ [13].

3. The Homotopy Inertia Group

In [15], the concordance inertia group $I_c(M^{2n})$ and the inertia group $I(M^{2n})$ of an $(n-1)$ -connected $2n$ -smooth manifold M^{2n} were considered for $n = 4, 8$. In this section, we compute the homotopy inertia group of M^{2n} .

Theorem 3.1. *Let $m \geq 5$ and M^m be a closed simply connected smooth m -manifold such that the integral and mod-2 cohomologies of M^m vanish in odd degrees. Then $F_{Con} : \mathcal{C}^{Diff}(M) \rightarrow \mathcal{S}^{Diff}(M)$ is injective.*

Proof. For $i \in \{1, 2\}$, let $[N_i, f_i] \in \mathcal{C}^{Diff}(M)$, where N_i is a closed smooth manifold and f_i is a homeomorphism from N_i to M such that $F_{Con}([N_1, f_1]) = F_{Con}([N_2, f_2])$. Thus, (N_1, f_1) and (N_2, f_2) represent the same element in $\mathcal{S}^{Diff}(M)$, and therefore, there exists a diffeomorphism ϕ from N_1 to N_2 such that $f_2 \circ \phi \simeq f_1$.

Since each (N_i, f_i) represents an element in $\mathcal{C}^{PL}(M)$ which is isomorphic to $H^3(M; \mathbb{Z}_2) = 0 = \{[M, Id]\}$, there is a PL -homeomorphism k_i from N_i to M such that $Id \circ k_i$ coincides up to topological concordance, and hence up to homotopy with f_i . Therefore, $k_2 \circ \phi \simeq k_1$, say under the homotopy H . Define the homotopy equivalence $\tilde{H} : N_1 \times I \rightarrow M \times I$ by $\tilde{H}(x, t) = (H(x, t), t)$, where $(x, t) \in N_1 \times I$. By [33, Theorem 1], \tilde{H} is homotopic to a PL homeomorphism (mod $N_1 \times \partial I$). It is because such a deformation is obstructed by cohomology classes in $H^*(N_1 \times (I, \partial I); P_*)$. Since N_1 can have nonzero cohomology groups in even dimensions and P_i is zero for odd i , these cohomology groups are all zero. Thus, $k_2 \circ \phi$ is topological concordant to k_1 and hence $Id \circ k_2 \circ \phi$ to $Id \circ k_1$, and $f_2 \circ \phi$ to f_1 . This completes the proof by noting that both (N_1, f_1) and (N_2, f_2) represent the same element in $\mathcal{C}^{Diff}(M)$. \square

As a corollary to the result above, we have the following.

Corollary 3.2. *Let $m \geq 5$ and M^m be a closed simply connected smooth m -manifold such that the integral and mod-2 cohomologies of M^m vanish in odd degrees. Then $I_c(M) = I_h(M)$.*

Remark 3.3. (1) If M^{2n} is a closed simply connected smooth $2n$ -manifold such that the integral cohomology of M^{2n} vanishes in odd degrees, by Corollary 3.2, $I_h(M^{2n}) = I_c(M^{2n})$.

(2) If M^{2n} is a closed smooth manifold homotopy equivalent to $\mathbb{C}\mathbf{P}^n$, then $I_c(M^{2n})=0$ for $n \leq 8$ [16]. Therefore, by Corollary 3.2, $I_h(M^{2n}) = 0$, $n \leq 8$.

By [15, Theorem 2.7((i), (iii))] and Corollary 3.2, we have the following.

Corollary 3.4. *Let M^{2n} be a closed smooth $(n-1)$ -connected $2n$ -manifold. Then*

(1) *If $n = 4$, then $I_h(M^{2n}) = 0$.*

(2) *If $n = 8$ and $H^n(M; \mathbb{Z}) \cong \mathbb{Z}$, then $I_h(M^{2n}) = 0$.*

Since the concordance inertia group is a homotopy invariant and by Corollary 3.2, we have the following result.

Corollary 3.5. *If M and N have the same homotopy type of simply connected closed smooth m -manifolds such that the integral and mod-2 cohomologies of M vanish in odd degrees, then their homotopy inertia groups are equal.*

4. Smooth Structures on 7-connected 16-manifolds

In [15, Theorem 2.7(ii)], it was shown that the group $\mathcal{C}(M^{2n})$ of concordance classes of smoothings of a closed smooth $(n-1)$ -connected $2n$ -manifold M^{2n} is isomorphic to the group of smooth homotopy spheres Θ_{2n} for $n = 4$. In this section, we compute the group $\mathcal{C}(M^{2n})$ for $n = 8$.

We denote $X^{2n} = \mathbb{S}^n \cup_g \mathbb{D}^{2n}$, where $g : \mathbb{S}^{2n-1} \rightarrow \mathbb{S}^n$ is an attaching map. Let $i_n : [\mathbb{S}^n; SPL] \rightarrow [\mathbb{S}^n; SG]$ and $\iota_*^* : [X, -] \rightarrow [\mathbb{S}^n, -]$ be the induced maps by the natural maps $i : SPL \rightarrow SG$ and $\iota : \mathbb{S}^n \rightarrow X$, respectively. Denote by $f_X : X \rightarrow \mathbb{S}^{2n}$, a degree one map.

Theorem 4.1. (i) $\iota_{PL}^* : [X^{16}, SPL] \rightarrow [\mathbb{S}^8, SPL]$ is surjective.

(ii) $[X^{16}, SG]$ is isomorphic to either $\mathbb{Z}_4 \oplus \mathbb{Z}_2$ or $\mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2$.

The proof of Theorem 4.1 requires one fact which we prove below:

Lemma 4.2. Let $n = 0 \pmod{4}$, $n > 0$. Then $i_n : [\mathbb{S}^n, SPL] \rightarrow [\mathbb{S}^n, SG]$ is an isomorphism.

Proof. Consider the homotopy exact sequence for the fibration $SPL \rightarrow SG \rightarrow G/PL$:

$$\begin{aligned} \cdots \rightarrow \pi_{n+1}(SG) \rightarrow \pi_{n+1}(G/PL) \rightarrow \pi_n(SPL) \\ \xrightarrow{i_n} \pi_n(SG) \rightarrow \pi_n(G/PL) \rightarrow \cdots \end{aligned} \quad (4.1)$$

and by using the fact $\pi_n(SG) = \pi_n^S$ is a finite group, $\pi_{n+1}(G/PL) = 0$ and $\pi_n(G/PL) = \mathbb{Z}$, we have that $i_n : [\mathbb{S}^n, SPL] \rightarrow [\mathbb{S}^n, SG]$ is an isomorphism.

□

Proof of Theorem 4.1. (i) There is a commutative diagram:

$$\begin{array}{ccccccccc} & & & & & & & & [\mathbb{S}^{16}, SPL] \\ & & & & & & & & \downarrow \cong \\ & & & & & & & & [\mathbb{S}^{16}, SG] \\ & & & & & & & & \downarrow \\ & & & & & & & & \mathbb{Z} \cong [\mathbb{S}^{16}, G/PL] \\ & & & & & & & & \downarrow \\ [\mathbb{S}^9, SPL] & \xrightarrow{S(g)^*} & [\mathbb{S}^{16}, SPL] & \xrightarrow{f_X^*} & [X^{16}, SPL] & \xrightarrow{\iota_{PL}^*} & [\mathbb{S}^8, SPL] & \xrightarrow{g^*} & [\mathbb{S}^{15}, SPL] \\ \downarrow i_9 & & \cong \downarrow i_{16} & & \downarrow & & \cong \downarrow i_8 & & \downarrow i_{15} \\ [\mathbb{S}^9, SG] & \xrightarrow{S(g)^*} & [\mathbb{S}^{16}, SG] & \xrightarrow{f_X^*} & [X^{16}, SG] & \xrightarrow{\iota_G^*} & [\mathbb{S}^8, SG] & \xrightarrow{g^*} & [\mathbb{S}^{15}, SG] \\ \downarrow & & & & & & & & \\ [\mathbb{S}^9, G/PL] = 0 & & & & & & & & \end{array} \quad (4.2)$$

in which, the homomorphism

$$i_n : \pi_n(SPL) \rightarrow \pi_n(SG)$$

is an isomorphism by Lemma 4.2, where $n = 8, 16$ and the rows and columns are part of the long exact sequences obtained from the cofiber sequence $\mathbb{S}^{15} \rightarrow \mathbb{S}^8 \rightarrow X$ and the fiber sequence $SPL \rightarrow SG \rightarrow G/PL$, respectively. From this commutative diagram, it is seen that

$$\iota_{PL}^* : [X^{16}, SPL] \rightarrow [\mathbb{S}^8, SPL]$$

is surjective if and only if

$$g^* : \pi_8(SG) \rightarrow \pi_{15}(SG)$$

is the zero homomorphism. To show that $g^* : \pi_8(SG) \rightarrow \pi_{15}(SG)$ is the zero homomorphism, we will use Toda's notation [35, p. 189] for special elements in the stable stems π_n^S . Recall that π_*^S equal the direct sum of the π_n^S is an anti-commutative graded ring with respect to composition as multiplication and $\pi_n(SG)$ can be identified with the n th stable homotopy group of sphere π_n^S ; i.e., $\pi_n(SG) = \pi_n^S$ for $n \geq 1$. This fact is used to identify the map

$$g^* : \pi_8(SG) \rightarrow \pi_{15}(SG)$$

with the map

$$(S^q(g))^* : \pi_{8+q}(\mathbb{S}^q) \rightarrow \pi_{15+q}(\mathbb{S}^q),$$

where $S^k(g)$ is the k th iterated suspension of g . Since $\pi_8^S = \mathbb{Z}_2(\bar{\nu}) \oplus \mathbb{Z}_2(\epsilon)$, where $\bar{\nu}$ and ϵ are the generators of the first and second components of π_8^S . Let $f : \mathbb{S}^{8+q} \rightarrow \mathbb{S}^q$ represent either $\bar{\nu}$ or ϵ . First note that the homotopy class

$$[S(g)] = aS(\sigma) + x \in \pi_7^S \cong \mathbb{Z}_{16} \oplus \mathbb{Z}_{15},$$

where $a \in \mathbb{Z}$, $S(\sigma) \in \mathbb{Z}_{16}$ is the generator represented by the suspension of the Hopf map $\sigma : \mathbb{S}^{15} \rightarrow \mathbb{S}^8$ and $x \in \mathbb{Z}_{15}$ has odd order. Using this together

with the fact that $\pi_8^S = \mathbb{Z}_2(\bar{\nu}) \oplus \mathbb{Z}_2(\epsilon)$ has order 4, we see that

$$\begin{aligned} f \circ S^q(g) &= f \circ S^{q-1}(aS(\sigma) + x) \\ &= a(f \circ S^{q-1}(S(\sigma))) + (f \circ S^{q-1}(x)) \\ &= a(f \circ S^{q-1}(S(\sigma))) = a(f \circ S^q(\sigma)). \end{aligned}$$

Since $S^q(\sigma)$ represents the element $S(\sigma)$ in π_7^S and $\bar{\nu} \circ S(\sigma) = \epsilon \circ S(\sigma) = 0$ in π_{15}^S [35, Theorem 14.1(iii), p. 190], we have $f \circ S^q(\sigma) = 0$ in π_{15}^S . This implies that $f \circ S^q(g) = 0$ in π_{15}^S . Therefore, the map $(S^q(g))^* : \pi_{8+q}(\mathbb{S}^q) \rightarrow \pi_{15+q}(\mathbb{S}^q)$ is the zero homomorphism and hence $g^* : \pi_8(SG) \rightarrow \pi_{15}(SG)$ is the zero homomorphism.

(ii) Recall the homotopy class

$$[S(g)] = a\sigma + x \in \pi_7^S \cong \mathbb{Z}_{16} \oplus \mathbb{Z}_{15},$$

where $a \in \mathbb{Z}$, σ itself denotes the generator of \mathbb{Z}_{16} represented by the suspension of the Hopf map $\sigma : \mathbb{S}^{15} \rightarrow \mathbb{S}^8$ and $x \in \mathbb{Z}_{15}$ has odd order. Using this together with the fact that $\pi_{16} = \mathbb{Z}_2(\eta^*) \oplus \mathbb{Z}_2(\eta \circ \rho)$ has order 4, we see that $\text{Im}(S(g)^* : [\mathbb{S}^9, SG] \rightarrow [\mathbb{S}^{16}, SG]) \subseteq \pi_{16}^S$ is generated by the following three elements:

$$\nu^3 \circ a\sigma, \quad \mu \circ a\sigma, \quad \eta \circ \epsilon \circ a\sigma,$$

where ν^3 , μ and $\eta \circ \epsilon$ are the generators of the first, second and third components of

$$\pi_9^S = [\mathbb{S}^9, SG] \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2,$$

respectively. Now, by [35, Theorem 14.1((i), (ii)), p. 190], we have

$$\nu \circ a\sigma = a(\nu \circ \sigma) = 0,$$

$$\sigma \circ \epsilon = 0$$

and $\sigma \circ \mu = \eta \circ \rho$. Using these relations and anti-commutativity of π_*^s , we have the following relations:

$$\begin{aligned} v^3 \circ a\sigma &= a(v^2 \circ (v \circ \sigma)) = 0, \\ \eta \circ \epsilon \circ a\sigma &= a(\eta \circ (\epsilon \circ \sigma)) = a(\eta \circ (\sigma \circ \epsilon)) = 0, \\ \mu \circ a\sigma &= a(\mu \circ \sigma) = -a(\sigma \circ \mu) = -a(\eta \circ \rho) = a(\rho \circ \mu). \end{aligned}$$

This implies that $\text{Im}(S(g)^* : [\mathbb{S}^9, SG] \rightarrow [\mathbb{S}^{16}, SG])$ is contained in the subgroup of π_{16}^s generated by $\rho \circ \mu$. This shows that $\text{Im}(S(g)^* : [\mathbb{S}^9, SG] \rightarrow [\mathbb{S}^{16}, SG]) = \mathbb{Z}_2$. Now, by the exact sequence in the horizontal rows of the commutative diagram (4.2) along SG and using the fact we have observed in the proof of (i) that $\iota_G^* : [X^{16}, SG] \rightarrow [\mathbb{S}^8, SG]$ is surjective, we get that

$$[X^{16}, SG] \cong \mathbb{Z}_4 \oplus \mathbb{Z}_2 \text{ or } \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2.$$

This completes the proof of Theorem 4.1. \square

The following result follows from Theorem 4.1.

Corollary 4.3. *Let M^{16} be a closed 7-connected 16-manifold such that $H^8(M; \mathbb{Z}) \cong \mathbb{Z}$. Then $[M, SG]$ is isomorphic to either $\mathbb{Z}_4 \oplus \mathbb{Z}_2$ or $\mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2$.*

Theorem 4.4. *Let M^8 be a closed 3-connected 8-manifold. Then*

- (i) $[M, SG] \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2$,
- (ii) $[M, G/O] \cong \left(\bigoplus_{i=1}^{k+1} \mathbb{Z} \right) \oplus \mathbb{Z}_2$,

where k is the 4th Betti number of M .

Proof. According to Wall [37], M has the homotopy type of $X = \left(\bigvee_{i=1}^k \mathbb{S}_i^4 \right) \cup_g \mathbb{D}^8$, where k is the 4th Betti number of M , $\bigvee_{i=1}^k \mathbb{S}_i^4$ is the

wedge sum of 4-spheres and $g : \mathbb{S}^7 \rightarrow \bigvee_{i=1}^k \mathbb{S}_i^4$ is the attaching map of \mathbb{D}^8 . Let $f_X : X \rightarrow \mathbb{S}^8$ be the collapsing map obtained by identifying \mathbb{S}^8 with $X/\bigvee_{i=1}^k \mathbb{S}_i^4$. Then by the naturality of the Puppe sequence, we have the following exact ladder:

$$\begin{aligned} \dots \rightarrow \left[\bigvee_{i=1}^k S\mathbb{S}_i^4, Cat \right] &\xrightarrow{(S(g))^*} [\mathbb{S}^8, Cat] \xrightarrow{f_X^*} [X, Cat] \\ \downarrow \iota_{Cat}^* & \\ \left[\bigvee_{i=1}^k \mathbb{S}_i^4, Cat \right] &\xrightarrow{g^*} [\mathbb{S}^7, Cat], \end{aligned} \tag{4.3}$$

where $S(g)$ is the suspension of the map $g : \mathbb{S}^7 \rightarrow \bigvee_{i=1}^k \mathbb{S}_i^4$.

Using the facts that

$$\left[\bigvee_{i=1}^k S\mathbb{S}_i^4, Cat \right] \cong \prod_{i=1}^k [\mathbb{S}_i^5, Cat]$$

and

$$\left[\bigvee_{i=1}^k \mathbb{S}_i^4, Cat \right] \cong \prod_{i=1}^k [\mathbb{S}_i^4, Cat],$$

the above exact sequence (4.3) becomes

$$\dots \rightarrow \prod_{i=1}^k [\mathbb{S}_i^5, Cat] \xrightarrow{(S(g))^*} [\mathbb{S}^8, Cat] \xrightarrow{f_X^*} [X, Cat] \xrightarrow{\iota_{Cat}^*} \prod_{i=1}^k [\mathbb{S}_i^4, Cat]. \tag{4.4}$$

If $Cat = SG$ in the exact sequence (4.4) and using the fact that $[\mathbb{S}^n, SG] = 0$, where $n = 4, 5$, we get that

$$[\mathbb{S}^8, SG] \cong [X, SG] \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2.$$

Hence

$$[M^8, SG] \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2.$$

This proves (i).

We now take $Cat = G/O$ and by using the facts that $[\mathbb{S}^8, G/O] = \mathbb{Z}_2 \oplus \mathbb{Z}$ and $[\mathbb{S}^n, G/O] = 0$ [33], where $n = 5, 7$, it follows that

$$[X, G/O] \cong [M, G/O] \cong \left(\bigoplus_{i=1}^{k+1} \mathbb{Z} \right) \oplus \mathbb{Z}_2.$$

This proves (ii). □

Recall that PL/O has a commutative H -space structure which makes the fibration $PL/O \rightarrow BSO \rightarrow BSPL$ into a sequence of H -space maps where BSO and $BSPL$ have compatible commutative H -space structures coming from the Whitney sum of bundles [23]. Associated to this fibration, we have the long exact Puppe sequences of abelian groups, for any space X ,

$$\cdots \rightarrow [X, SPL] \xrightarrow{\phi_*} [X, PL/O] \xrightarrow{\partial_X} [X, BSO] \rightarrow [X, BSPL], \quad (4.5)$$

where ϕ_* is induced by the natural map $\phi : SPL \rightarrow PL/O$. When $X = M$ is a smooth manifold, ∂_M computes the difference a smooth structure makes to the isomorphism class of the stable tangent bundle. That is,

$$\partial_M([N, f]) = (f^{-1})^* T^0(N) - T^0(M) \in \widetilde{KO}^0(M) = [M, BSO],$$

where $T^0(M)$ is the stable tangent bundle of M .

We now prove the following.

Theorem 4.5. *Let M^{16} be a closed smooth 7-connected 16-manifold such that $H^8(M; \mathbb{Z}) \cong \mathbb{Z}$. Then $\mathcal{C}^{Diff}(M)$ contains exactly four elements.*

Proof. Since PL/O fits into a fibration

$$PL/O \rightarrow Top/O \rightarrow Top/PL \simeq K(\mathbb{Z}_2, 3),$$

we have the following long exact sequence:

$$\begin{aligned}
 \cdots [M, \Omega(\text{Top}/PL)] &= [M, K(\mathbb{Z}_2, 2)] = H^2(M; \mathbb{Z}_2) \\
 &= 0 \rightarrow [M, PL/O] \rightarrow [M, \text{Top}/O] \rightarrow [M, \text{Top}/PL] \\
 &= [M, K(\mathbb{Z}_2, 3)] = H^3(M; \mathbb{Z}_2) = 0.
 \end{aligned}$$

This implies that $[M, PL/O] \cong [M, \text{Top}/O]$. To show that $\mathcal{C}^{\text{Diff}}(M)$ contains exactly four elements, it is enough to show that $[M, PL/O]$ contains exactly four elements by using the identifications $\mathcal{C}^{\text{Diff}}(M^{16}) = [M, \text{Top}/O]$ given by [22, pp. 194-196]. Since M^{16} is a closed 7-connected 16-manifold such that $H^8(M; \mathbb{Z}) \cong \mathbb{Z}$, M has the homotopy type of $X = \mathbb{S}^8 \cup_{\alpha} \mathbb{D}^{16}$, where $\alpha : \mathbb{S}^{15} \rightarrow \mathbb{S}^8$ is the attaching map of \mathbb{D}^{16} .

Consider the following commuting diagram of long exact sequences:

$$\begin{array}{ccccccc}
 \cdots & \longrightarrow & [X, SPL] & \xrightarrow{\phi_*} & [X, PL/O] & \xrightarrow{\partial_X} & [X, BSO] \longrightarrow \cdots \\
 & & \downarrow \iota^* & & \downarrow \iota^* & & \downarrow \iota^* \\
 \cdots & \longrightarrow & [\mathbb{S}^8, SPL] & \xrightarrow{\phi_*} & [\mathbb{S}^8, PL/O] = \mathbb{Z}_2 & \xrightarrow{\partial_{\mathbb{S}^8}} & [\mathbb{S}^8, BSO] \longrightarrow \cdots
 \end{array} \quad (4.6)$$

and using the facts that $\phi_* : [\mathbb{S}^8, SPL] \rightarrow [\mathbb{S}^8, PL/O]$ and $\iota^* : [X, SPL] \rightarrow [\mathbb{S}^8, SPL]$ are surjective by the result of [6] and Theorem 4.1, respectively, we have that $\iota^* : [X, PL/O] \rightarrow [\mathbb{S}^8, PL/O]$ is surjective. Since $[\mathbb{S}^8, PL/O] \cong \mathbb{Z}_2$, the homomorphism $\iota^* : [X, PL/O] \rightarrow [\mathbb{S}^8, PL/O]$ fits into the following long exact Puppe sequences of abelian groups:

$$\cdots \rightarrow [\mathbb{S}^{16}, PL/O] \xrightarrow{f_X^*} [X, PL/O] \xrightarrow{\iota^*} [\mathbb{S}^8, PL/O] \xrightarrow{\sigma} [\mathbb{S}^{15}, PL/O], \quad (4.7)$$

where $f_X^* : [\mathbb{S}^{16}, PL/O] = \Theta_{16} \rightarrow [X, PL/O]$ is monic by [15, Theorem 2.7(iii)]. This implies that $[X, PL/O] \cong [M, PL/O]$ contains exactly four elements. This proves the theorem. \square

In [15], it was proved that there are, up to concordance, exactly two smooth structures on a closed smooth 3-connected 8-manifold. Now, Theorem 4.5 implies the following.

Corollary 4.6. *Let M^{16} be a closed smooth 7-connected 16-manifold such that $H^8(M; \mathbb{Z}) \cong \mathbb{Z}$. Then M^{16} has exactly four distinct smooth structures up to concordance.*

5. The Smooth Tangential Structure Set

In order to determine the smooth tangential structure set $\mathcal{S}^{TDiff}(M)$, one must compute the surgery obstruction map σ . We now prove the following.

Theorem 5.1. *Let M^{2n} be a closed smooth $(n-1)$ -connected $2n$ -manifold with $n \equiv 0 \pmod{4}$, $n > 0$. Then the surgery obstruction $\sigma : [M^{2n}, SG] \rightarrow L_{2n}(\mathbb{Z}) \cong \mathbb{Z}$ is the zero map.*

Let $\phi_* : [M, SG] \rightarrow [M, G/O]$ and $\psi_* : [M, Top/O] \rightarrow [M, G/O]$ be the induced maps by the natural maps $\phi : SG \rightarrow G/O$ and $\psi : Top/O \rightarrow G/O$, respectively. We prove the following.

Theorem 5.2. *Let M^{2n} be a closed smooth $(n-1)$ -connected $2n$ -manifold with $n \equiv 0 \pmod{4}$, $n > 0$. Then*

$$\begin{aligned} & \text{Im}(\phi_* : [M^{2n}, SG] \rightarrow [M^{2n}, G/O]) \\ &= \text{Im}(\psi_* : [M^{2n}, Top/O] \rightarrow [M^{2n}, G/O]) \cong [M^{2n}, Top/O]. \end{aligned}$$

Proof. Consider the following commutative diagram:

$$\begin{array}{ccccc}
 & & [M^{2n}, \Omega(G/Top)] & & \\
 & & \downarrow & & \\
 & & [M^{2n}, Top/O] & \equiv & [M^{2n}, Top/O] \\
 & & \downarrow \psi_* & & \downarrow \partial_M \\
 [M^{2n}, SG] & \xrightarrow{\phi_*} & [M^{2n}, G/O] & \xrightarrow{\partial_M} & [M^{2n}, BSO] \\
 \parallel & & \downarrow & & \downarrow \\
 [M^{2n}, SG] & \xrightarrow{\phi_*} & [M^{2n}, G/Top] & \xrightarrow{\partial_M} & [M^{2n}, BStop]
 \end{array} \tag{5.1}$$

In this diagram, the rows and columns are the parts of the long exact sequences obtained from the fiber sequences $SG \rightarrow G/O \rightarrow BSO$, $SG \rightarrow G/Top \rightarrow BStop$ and $Top/O \rightarrow G/O \rightarrow G/Top$.

Now the following three facts used in conjunction with a simple diagram chase in (5.1) show that

$$\begin{aligned}
 & \text{Im}(\phi_* : [M^{2n}, SG] \rightarrow [M^{2n}, G/O]) \\
 &= \text{Im}(\psi_* : [M^{2n}, Top/O] \rightarrow [M^{2n}, G/O]) \cong [M^{2n}, Top/O].
 \end{aligned}$$

thus proving Theorem 5.2. \square

Fact 1. The homomorphism $\partial_M : [M^{2n}, G/Top] \rightarrow [M^{2n}, BStop]$ is monic for $n \equiv 0 \pmod{4}$.

Fact 2. The homomorphism $\psi_* : [M^{2n}, Top/O] \rightarrow [M^{2n}, G/O]$ is monic.

Fact 3. The homomorphism $\partial_M : [M^{2n}, Top/O] \rightarrow [M^{2n}, BSO]$ is the zero map.

It remains to verify these facts. Fact 1 is due to Kramer [21, Proposition 9.9].

Proof of Fact 2. Consider the Puppe's exact sequence (4.4) for the inclusion $i : \bigvee_{i=1}^{i=k} \mathbb{S}_i^n \rightarrow M^{2n}$ along $Cat = \Omega(G/Top)$ and using the fact that $[\mathbb{S}^n, \Omega(G/Top)] = 0$ for even n , we get that $[M^{2n}, \Omega(G/Top)] = 0$. Now it follows from the vertical exact sequence in (5.1) that the homomorphism $\psi_* : [M^{2n}, Top/O] \rightarrow [M^{2n}, G/O]$ is monic. \square

Proof of Fact 3. Since $[M^{2n}, Top/O]$ is a finite group and $\widetilde{KO}(M^{2n}) = [M^{2n}, BSO]$ [41] is torsion-free, it follows that the homomorphism $\partial_M : [M^{2n}, Top/O] \rightarrow [M^{2n}, BSO]$ is the zero map. \square

Proof of Theorem 5.1. The smooth structure set $\mathcal{S}^{Diff}(M^{2n})$ fits into the following surgery exact sequence:

$$\mathcal{S}^{Diff}(M) \xrightarrow{inj} [M, G/O] \xrightarrow{\sigma} L_{2n}(\mathbb{Z}). \quad (5.2)$$

Thus, a normal map $\varphi \in [M, G/O]$ represents a homotopy smoothing structure for M if and only if $\sigma(\varphi) = 0$. Observe that if φ lies in the image $[M^{2n}, Top/O] \xrightarrow{\psi_*} \mathcal{S}^{Diff}(M^{2n}) \xrightarrow{inj} [M^{2n}, G/O]$, then $\sigma(\varphi) = 0$. Therefore, the surgery obstruction $\sigma : \text{Im}(\psi_* : [M^{2n}, Top/O] \rightarrow [M^{2n}, G/O]) \rightarrow L_{2n}(\mathbb{Z})$ is the zero map. Combining this with the following commutative diagram [9] and Theorem 5.2:

$$\begin{array}{ccc} [M^{2n}, SG] & \xrightarrow{\sigma} & L_{2n}(\mathbb{Z}) \\ \phi_* \downarrow & & \parallel \\ [M^{2n}, G/O] & \xrightarrow{\sigma} & L_{2n}(\mathbb{Z}) \end{array} \quad (5.3)$$

we get that the surgery obstruction $\sigma : [M^{2n}, SG] \rightarrow L_{2n}(\mathbb{Z}) \cong \mathbb{Z}$ is the zero map. This completes the proof of the theorem. \square

Theorem 5.3. *Let M^{2n} be a closed smooth $(n-1)$ -connected $2n$ -manifold with $n \equiv 0 \pmod{4}$, $n > 0$. Then the map $\eta^t : \mathcal{S}^{TDiff}(M^{2n}) \rightarrow [M^{2n}, SG]$ is a set bijection.*

Proof. Since $L_{2n+1}(\mathbb{Z}) = 0$, the tangential surgery exact sequence for M^{2n} runs as follows:

$$0 \rightarrow \mathcal{S}^{TDiff}(M^{2n}) \xrightarrow{\eta^t} [M^{2n}, SG] \xrightarrow{\sigma} L_{2n}(\mathbb{Z}). \quad (5.4)$$

From Theorem 5.1, it follows that the map $\eta^t : \mathcal{S}^{TDiff}(M^{2n}) \rightarrow [M^{2n}, SG]$ is a set bijection. \square

Theorem 5.4. *Let M^{2n} be a closed smooth $(n-1)$ -connected $2n$ -manifold with $n \equiv 0 \pmod{4}$, $n > 0$. Let $f : N \rightarrow M$ be a tangential homotopy equivalence where N is a closed smooth manifold. Then there exist a smooth manifold \tilde{N} and a homeomorphism $g : \tilde{N} \rightarrow M$ such that (N, f) and (\tilde{N}, g) represent the same element in $\mathcal{S}^{Diff}(M)$.*

Proof. Observe that a tangential homotopy equivalence $f : N \rightarrow M$ represents an element of $\mathcal{S}^{Diff}(M)$ and its image in $\widetilde{KO}^0(M) = [M, BSO]$ is given by $(f^{-1})^*T^0(N) - T^0(M) = 0$. Therefore, by using the third row exact sequence in the diagram (5.1) and Theorem 5.2, there exist a smooth manifold \tilde{N} and a homeomorphism $g : \tilde{N} \rightarrow M$ such that (N, f) and (\tilde{N}, g) represent the same map in $[M, G/O]$. Thus, (N, f) and (\tilde{N}, g) represent the same element in $\mathcal{S}^{Diff}(M)$. \square

We note that Theorem 5.4 implies the following.

Corollary 5.5. *Let M^{2n} be a closed smooth $(n-1)$ -connected $2n$ -manifold with $n \equiv 0 \pmod{4}$, $n > 0$. If a smooth manifold N is tangential homotopy equivalent to M , then N is homeomorphic to M .*

Theorem 5.6. *Let M be a closed smooth 3-connected 8-manifold. If a smooth manifold N is tangential homotopy equivalent to M , then there is a homotopy 8-sphere Σ such that N is diffeomorphic to $M \# \Sigma$.*

Proof. By [15, Theorem 2.7(ii)], we have $\mathcal{C}^{Diff}(M) = \{[M\#\Sigma] \mid \Sigma \in \Theta_8\}$. Now applying Theorem 5.4. \square

As a consequence of [15, Theorem 1.1] and Theorem 5.6, we get

Corollary 5.7. *Let M be a closed smooth 3-connected 8-manifold such that $H^4(M; \mathbb{Z}) \cong \mathbb{Z}$. If a smooth manifold N is tangential homotopy equivalent to M , then N is diffeomorphic to M .*

Acknowledgement

The author thanks the anonymous referees for their valuable suggestions which led to the improvement of the manuscript.

References

- [1] D. W. Anderson, E. H. Brown and F. P. Peterson, The structure of the spin cobordism ring, *Ann. of Math. (2)* 86 (1967), 271-298.
- [2] C. S. Aravinda and F. T. Farrell, Exotic negatively curved structures on Cayley hyperbolic manifolds, *J. Differential Geom.* 63 (2003), 41-62.
- [3] D. Barden, Simply connected five-manifolds, *Ann. of Math. (2)* 82 (1965), 365-385.
- [4] W. Browder, *Surgery on simply-connected manifolds*, *Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 65*, Springer-Verlag, New York, Heidelberg, 1972.
- [5] G. Brumfiel, Homotopy equivalences of almost smooth manifolds, *Comment. Math. Helv.* 46 (1971), 381-407.
- [6] G. Brumfiel, On the homotopy groups of BPL and PL/O , *Ann. of Math. (2)* 88 (1968), 291-311.
- [7] D. Crowley, The smooth structure set of $\mathbb{S}^p \times \mathbb{S}^q$, *Geom. Dedicata* 148 (2010), 15-33.
- [8] D. Crowley, On the classification of highly connected manifolds in dimensions 7 and 15, Ph.D. Thesis, 2002.
- [9] D. Crowley and I. Hambleton, Finite group actions on Kervaire manifolds, *Adv. Math.* 283 (2015), 88-129.

- [10] D. Crowley and J. Nordström, The classification of 2-connected 7-manifolds, 2014, preprint.
- [11] J. Eells, Jr. and N. H. Kuiper, Manifolds which are like projective planes, *Inst. Hautes Etudes Sci. Publ. Math.* 14 (1962), 5-46.
- [12] M. Frame, On the inertia groups of h -cobordant manifolds, *Proc. Amer. Math. Soc.* 90(3) (1984), 433-439.
- [13] B. Harris, J -homomorphisms and cobordism groups, *Invent. Math.* 7 (1969), 313-320.
- [14] M. Joachim and D. J. Wraith, Exotic spheres and curvature, *Bull. Amer. Math. Soc. (N.S.)* 45(4) (2008), 595-616.
- [15] R. Kasilingam, Inertia groups and smooth structures of $(n - 1)$ -connected $2n$ -manifolds, *Osaka J. Math.* 53(2) (2016), 309-319.
- [16] K. Kawakubo, Inertia groups of low dimensional complex projective spaces and some free differentiable actions on spheres I, *Proc. Japan Acad.* 44 (1968), 873-875.
- [17] K. Kawakubo, On the inertia groups of homology tori, *J. Math. Soc. Japan* 21 (1969), 37-47.
- [18] M. A. Kervaire and J. W. Milnor, Groups of homotopy spheres I, *Ann. of Math. (2)* 77 (1963), 504-537.
- [19] A. Kosinski, On the inertia group of π -manifolds, *Amer. J. Math.* 89 (1967), 227-248.
- [20] M. Kreck, Surgery and duality, *Ann. of Math. (2)* 149(3) (1999), 707-754.
- [21] L. Kramer, Projective planes and their look-alikes, *J. Differential Geom.* 64(1) (2003), 1-55.
- [22] R. C. Kirby and L. C. Siebenmann, Foundational essays on topological manifolds, smoothings, and triangulations, *Ann. of Math. Stud.* 88 (1977).
- [23] T. Lance, Differentiable structures on manifolds, *Surveys on Surgery Theory*, Vol. 1, *Ann. of Math. Stud.*, 145, Princeton Univ. Press, Princeton, NJ, 2000, pp. 73-104.
- [24] W. Luck, A basic introduction to surgery theory, *ICTP Lecture Notes Series* 9, Band 1, of the School High-dimensional Manifold Theory in Trieste, May/June 2001, Abdus Salam International Centre for Theoretical Physics, Trieste, 2002.
- [25] J. W. Milnor, On manifolds homeomorphic to the 7-sphere, *Ann. of Math. (2)* 64 (1956), 399-405.

- [26] J. Milnor, Some consequences of a theorem of Bott, *Ann. of Math. (2)* 68 (1958), 444-449.
- [27] I. Madsen, L. R. Taylor and B. Williams, Tangential homotopy equivalences, *Comment. Math. Helv.* 55 (1980), 445-484.
- [28] A. A. Ranicki, Algebraic L -theory and topological manifolds, *Cambridge Tracts in Math.*, Vol. 102, Cambridge Univ. Press, Cambridge, 1992.
- [29] A. A. Ranicki, Algebraic and geometric surgery, *Oxford Mathematical Monographs*, Oxford University Press, 2002.
- [30] R. Schultz, On the inertia group of a product of spheres, *Trans. Amer. Math. Soc.* 156 (1971), 137-153.
- [31] S. Smale, On the structure of manifolds, *Amer. J. Math.* 84 (1962), 387-399.
- [32] R. E. Stong, *Notes on Cobordism Theory*, Princeton Univ. Press, Princeton, N.J., 1968.
- [33] D. Sullivan, Triangulating and smoothing homotopy equivalences and homeomorphisms, *The Hauptvermutung Book, A Collection of Papers of the Topology of Manifolds, K-Monogr. Math.* 1, Kluwer Academic Publishers, Dordrecht, 1996, pp. 69-103.
- [34] I. Tamura, 8-manifolds admitting no differentiable structure, *J. Math. Soc. Japan* 13 (1961), 377-382.
- [35] H. Toda, Composition methods in homotopy groups of spheres, *Ann. of Math. Stud.*, No. 49, Princeton University Press, Princeton, N.J., 1962.
- [36] A. Tsuchiya, Characteristic classes for spherical fibre spaces, *Nagoya Math. J.* 43 (1971), 1-39.
- [37] C. T. C. Wall, Classification of $(n - 1)$ -connected $2n$ -manifolds, *Ann. of Math. (2)* 75 (1962), 163-189.
- [38] C. T. C. Wall, *Surgery on compact manifolds*, Second Edition, Edited and with a foreword by A. A. Ranicki, *Mathematical Surveys and Monographs*, 69, American Mathematical Society, Providence, RI, 1999.
- [39] S. Weinberger, On smooth surgery, *Comm. Pure Appl. Math.* 43(5) (1990), 695-696.
- [40] G. W. Whitehead, *Elements of homotopy theory*, *Graduate Texts in Math.*, Vol. 61, Springer-Verlag, New York, Heidelberg, Berlin, 1978.
- [41] H. Yang, Almost complex structures on $(n - 1)$ -connected $2n$ -manifolds, *Topology Appl.* 159(5) (2011), 1361-1368.