

## INERTIA GROUPS AND SMOOTH STRUCTURES OF $(n - 1)$ -CONNECTED $2n$ -MANIFOLDS

KASILINGAM RAMESH

(Received July 2, 2014, revised December 22, 2014)

### Abstract

Let  $M^{2n}$  denote a closed  $(n - 1)$ -connected smoothable topological  $2n$ -manifold. We show that the group  $\mathcal{C}(M^{2n})$  of concordance classes of smoothings of  $M^{2n}$  is isomorphic to the group of smooth homotopy spheres  $\bar{\Theta}_{2n}$  for  $n = 4$  or  $5$ , the concordance inertia group  $I_c(M^{2n}) = 0$  for  $n = 3, 4, 5$  or  $11$  and the homotopy inertia group  $I_h(M^{2n}) = 0$  for  $n = 4$ . On the way, following Wall's approach [16] we present a new proof of the main result in [9], namely, for  $n = 4, 8$  and  $H^n(M^{2n}; \mathbb{Z}) \cong \mathbb{Z}$ , the inertia group  $I(M^{2n}) \cong \mathbb{Z}_2$ . We also show that, up to orientation-preserving diffeomorphism,  $M^8$  has at most two distinct smooth structures;  $M^{10}$  has exactly six distinct smooth structures and then show that if  $M^{14}$  is a  $\pi$ -manifold,  $M^{14}$  has exactly two distinct smooth structures.

### 1. Introduction

We work in the categories of closed, oriented, simply-connected  $Cat$ -manifolds  $M$  and  $N$  and orientation preserving maps, where  $Cat = Diff$  for smooth manifolds or  $Cat = Top$  for topological manifolds. Let  $\bar{\Theta}_m$  be the group of smooth homotopy spheres defined by M. Kervaire and J. Milnor in [6]. Recall that the collection of homotopy spheres  $\Sigma$  which admit a diffeomorphism  $M \rightarrow M \# \Sigma$  form a subgroup  $I(M)$  of  $\bar{\Theta}_m$ , called the inertia group of  $M$ , where we regard the connected sum  $M \# \Sigma^m$  as a smooth manifold with the same underlying topological space as  $M$  and with smooth structure differing from that of  $M$  only on an  $m$ -disc. The homotopy inertia group  $I_h(M)$  of  $M^m$  is a subset of the inertia group consisting of homotopy spheres  $\Sigma$  for which the identity map  $\text{id}: M \rightarrow M \# \Sigma^m$  is homotopic to a diffeomorphism. Similarly, the concordance inertia group of  $M^m$ ,  $I_c(M^m) \subseteq \bar{\Theta}_m$ , consists of those homotopy spheres  $\Sigma^m$  such that  $M$  and  $M \# \Sigma^m$  are concordant.

The paper is organized as following. Let  $M^{2n}$  denote a closed  $(n - 1)$ -connected smoothable topological  $2n$ -manifold. In Section 2, we show that the group  $\mathcal{C}(M^{2n})$  of concordance classes of smoothings of  $M^{2n}$  is isomorphic to the group of smooth homotopy spheres  $\bar{\Theta}_{2n}$  for  $n = 4$  or  $5$ , the concordance inertia group  $I_c(M^{2n}) = 0$  for  $n = 3, 4, 5$  or  $11$  and the homotopy inertia group  $I_h(M^{2n}) = 0$  for  $n = 4$ .

In Section 3, we present a new proof of the following result in [9].

**Theorem 1.1.** *Let  $M^{2n}$  be an  $(n - 1)$ -connected closed smooth manifold of dimension  $2n \neq 4$  such that  $H^n(M; \mathbb{Z}) \cong \mathbb{Z}$ . Then the inertia group  $I(M^{2n}) \cong \mathbb{Z}_2$ .*

In Section 4, we show that, up to orientation-preserving diffeomorphism,  $M^8$  has at most two distinct smooth structures;  $M^{10}$  has exactly six distinct smooth structures and if  $M^{14}$  is a  $\pi$ -manifold, then  $M^{14}$  has exactly two distinct smooth structures.

## 2. Concordance inertia groups of $(n - 1)$ -connected $2n$ -manifolds

We recall some terminology from [6]:

DEFINITION 2.1. (a) A homotopy  $m$ -sphere  $\Sigma^m$  is a closed oriented smooth manifold homotopy equivalent to the standard unit sphere  $\mathbb{S}^m$  in  $\mathbb{R}^{m+1}$ .

(b) A homotopy  $m$ -sphere  $\Sigma^m$  is said to be exotic if it is not diffeomorphic to  $\mathbb{S}^m$ .

DEFINITION 2.2. Define the  $m$ -th group of smooth homotopy spheres  $\Theta_m$  as follows. Elements are oriented  $h$ -cobordism classes  $[\Sigma]$  of homotopy  $m$ -spheres  $\Sigma$ , where  $\Sigma$  and  $\Sigma'$  are called (oriented)  $h$ -cobordant if there is an oriented  $h$ -cobordism  $(W, \partial_0 W, \partial_1 W)$  together with orientation preserving diffeomorphisms  $\Sigma \rightarrow \partial_0 W$  and  $(\Sigma')^- \rightarrow \partial_1 W$ . The addition is given by the connected sum. The zero element is represented by  $\mathbb{S}^m$ . The inverse of  $[\Sigma]$  is given by  $[\Sigma^-]$ , where  $\Sigma^-$  is obtained from  $\Sigma$  by reversing the orientation. M. Kervaire and J. Milnor [6] showed that each  $\Theta_m$  is a finite abelian group ( $m \geq 1$ ).

DEFINITION 2.3. Two homotopy  $m$ -spheres  $\Sigma_1^m$  and  $\Sigma_2^m$  are said to be equivalent if there exists an orientation preserving diffeomorphism  $f: \Sigma_1^m \rightarrow \Sigma_2^m$ .

The set of equivalence classes of homotopy  $m$ -spheres is denoted by  $\bar{\Theta}_m$ . The Kervaire–Milnor [6] paper worked rather with the group  $\Theta_m$  of smooth homotopy spheres up to  $h$ -cobordism. This makes a difference only for  $m = 4$ , since it is known, using the  $h$ -cobordism theorem of Smale [12], that  $\Theta_m \cong \bar{\Theta}_m$  for  $m \neq 4$ . However the difference is important in the four dimensional case, since  $\Theta_4$  is trivial, while the structure of  $\bar{\Theta}_4$  is a great unsolved problem.

DEFINITION 2.4. Let  $M$  be a closed topological manifold. Let  $(N, f)$  be a pair consisting of a smooth manifold  $N$  together with a homeomorphism  $f: N \rightarrow M$ . Two such pairs  $(N_1, f_1)$  and  $(N_2, f_2)$  are concordant provided there exists a diffeomorphism  $g: N_1 \rightarrow N_2$  such that the composition  $f_2 \circ g$  is topologically concordant to  $f_1$ , i.e., there exists a homeomorphism  $F: N_1 \times [0, 1] \rightarrow M \times [0, 1]$  such that  $F|_{N_1 \times 0} = f_1$  and  $F|_{N_1 \times 1} = f_2 \circ g$ . The set of all such concordance classes is denoted by  $\mathcal{C}(M)$ .

We will denote the class in  $\mathcal{C}(M)$  of  $(M^m \# \Sigma^m, \text{id})$  by  $[M^m \# \Sigma^m]$ . (Note that  $[M^n \# \mathbb{S}^n]$  is the class of  $(M^n, \text{id})$ .)

**DEFINITION 2.5.** Let  $M^m$  be a closed smooth  $m$ -dimensional manifold. The inertia group  $I(M) \subset \bar{\Theta}_m$  is defined as the set of  $\Sigma \in \bar{\Theta}_m$  for which there exists a diffeomorphism  $\phi: M \rightarrow M \# \Sigma$ .

Define the homotopy inertia group  $I_h(M)$  to be the set of all  $\Sigma \in I(M)$  such that there exists a diffeomorphism  $M \rightarrow M \# \Sigma$  which is homotopic to  $\text{id}: M \rightarrow M \# \Sigma$ .

Define the concordance inertia group  $I_c(M)$  to be the set of all  $\Sigma \in I_h(M)$  such that  $M \# \Sigma$  is concordant to  $M$ .

- REMARK 2.6.** (1) Clearly,  $I_c(M) \subseteq I_h(M) \subseteq I(M)$ .  
 (2) For  $M = \mathbb{S}^m$ ,  $I_c(M) = I_h(M) = I(M) = 0$ .

Now we have the following:

**Theorem 2.7.** Let  $M^{2n}$  be a closed smooth  $(n - 1)$ -connected  $2n$ -manifold with  $n \geq 3$ .

- (i) If  $n$  is any integer such that  $\Theta_{n+1}$  is trivial, then  $I_c(M^{2n}) = 0$ .  
 (ii) If  $n$  is any integer greater than 3 such that  $\Theta_n$  and  $\Theta_{n+1}$  are trivial, then

$$\mathcal{C}(M^{2n}) = \{[M^{2n} \# \Sigma] \mid \Sigma \in \bar{\Theta}_{2n}\} \cong \bar{\Theta}_{2n}.$$

- (iii) If  $n = 8$  and  $H^n(M; \mathbb{Z}) \cong \mathbb{Z}$ , then  $M^{2n} \# \Sigma^{2n}$  is not concordant to  $M^{2n}$ , where  $\Sigma^{2n} \in \bar{\Theta}_{2n}$  is the exotic sphere. In particular,  $\mathcal{C}(M^{2n})$  has at least two elements.  
 (iv) If  $n$  is any even integer such that  $\Theta_n$  and  $\Theta_{n+1}$  are trivial, then  $I_h(M) = 0$ .

**Proof.** Let  $Cat = Top$  or  $G$ , where  $Top$  and  $G$  are the stable spaces of self homeomorphisms of  $\mathbb{R}^n$  and self homotopy equivalences of  $\mathbb{S}^{n-1}$  respectively. For any degree one map  $f_M: M \rightarrow \mathbb{S}^{2n}$ , we have a homomorphism

$$f_M^*: [\mathbb{S}^{2n}, Cat/O] \rightarrow [M, Cat/O].$$

By Wall [15],  $M$  has the homotopy type of  $X = (\bigvee_{i=1}^k \mathbb{S}_i^n) \cup_g \mathbb{D}^{2n}$ , where  $k$  is the  $n$ -th Betti number of  $M$ ,  $\bigvee_{i=1}^k \mathbb{S}_i^n$  is the wedge sum of  $n$ -spheres which is the  $n$ -skeleton of  $M$  and  $g: \mathbb{S}^{2n-1} \rightarrow \bigvee_{i=1}^k \mathbb{S}_i^n$  is the attaching map of  $\mathbb{D}^{2n}$ . Let  $\phi: M \rightarrow X$  be a homotopy equivalence of degree one and  $q: X \rightarrow \mathbb{S}^{2n}$  be the collapsing map obtained by identifying  $\mathbb{S}^{2n}$  with  $X/\bigvee_{i=1}^k \mathbb{S}_i^n$  in an orientation preserving way. Let  $f_M = q \circ \phi: M \rightarrow \mathbb{S}^{2n}$  be the degree one map.

Consider the following Puppe's exact sequence for the inclusion  $i: \bigvee_{i=1}^k \mathbb{S}_i^n \hookrightarrow X$  along  $Cat/O$ :

(2.1)

$$\dots \rightarrow \left[ \bigvee_{i=1}^k S\mathbb{S}_i^n, Cat/O \right] \xrightarrow{(S(g))^*} [\mathbb{S}^{2n}, Cat/O] \xrightarrow{q^*} [X, Cat/O] \xrightarrow{i^*} \left[ \bigvee_{i=1}^k \mathbb{S}_i^n, Cat/O \right],$$

where  $S(g)$  is the suspension of the map  $g: \mathbb{S}^{2n-1} \rightarrow \bigvee_{i=1}^k \mathbb{S}_i^n$ .

Using the fact that

$$\left[ \bigvee_{i=1}^k S\mathbb{S}_i^n, \text{Cat}/O \right] \cong \prod_{i=1}^k [\mathbb{S}_i^{n+1}, \text{Cat}/O]$$

and

$$\left[ \bigvee_{i=1}^k \mathbb{S}_i^n, \text{Cat}/O \right] \cong \prod_{i=1}^k [\mathbb{S}_i^n, \text{Cat}/O],$$

the above exact sequence (2.1) becomes

$$\dots \rightarrow \prod_{i=1}^k [\mathbb{S}_i^{n+1}, \text{Cat}/O] \xrightarrow{(S(g))^*} [\mathbb{S}^{2n}, \text{Cat}/O] \xrightarrow{q^*} [X, \text{Cat}/O] \xrightarrow{i^*} \prod_{i=1}^k [\mathbb{S}_i^n, \text{Cat}/O].$$

(i): If  $n$  is any integer such that  $\Theta_{n+1}$  is trivial and  $\text{Cat} = \text{Top}$  in the above exact sequence (2.1), by using the fact that

$$[\mathbb{S}^m, \text{Top}/O] = \bar{\Theta}_m \quad (m \neq 3, 4)$$

and  $[\mathbb{S}^4, \text{Top}/O] = 0$  ([10, pp.200–201]), we have  $q^*: [\mathbb{S}^{2n}, \text{Top}/O] \rightarrow [X, \text{Top}/O]$  is injective. Hence  $f_M^* = \phi^* \circ q^*: \bar{\Theta}_{2n} \rightarrow [M, \text{Top}/O]$  is injective. By using the identifications  $\mathcal{C}(M^{2n}) = [M, \text{Top}/O]$  given by [10, pp.194–196],  $f_M^*: \bar{\Theta}_{2n} \rightarrow \mathcal{C}(M^{2n})$  becomes  $[\Sigma^{2n}] \rightarrow [M \# \Sigma^{2n}]$ .  $I_c(M)$  is exactly the kernel of  $f_M^*$ , and so  $I_c(M) = 0$ . This proves (i).

(ii): If  $n > 3$ ,  $\Theta_n$  and  $\Theta_{n+1}$  are trivial, and  $\text{Cat} = \text{Top}$  then, from the above exact sequence (2.1) we have  $q^*: [\mathbb{S}^{2n}, \text{Top}/O] \rightarrow [X, \text{Top}/O]$  is an isomorphism. This shows that  $f_M^* = \phi^* \circ q^*: \bar{\Theta}_{2n} \rightarrow \mathcal{C}(M^{2n})$  is an isomorphism and hence

$$\mathcal{C}(M^{2n}) = \{[M^{2n} \# \Sigma] \mid \Sigma \in \bar{\Theta}_{2n}\}.$$

This proves (ii).

(iii): If  $n = 8$  and  $H^n(M; \mathbb{Z}) \cong \mathbb{Z}$ , then  $M^{2n}$  has the homotopy type of  $X = \mathbb{S}^n \cup_g \mathbb{D}^{2n}$ , where  $g: \mathbb{S}^{2n-1} \rightarrow \mathbb{S}^n$  is the attaching map. In order to prove  $M^{2n} \# \Sigma^{2n}$  is not concordant to  $M^{2n}$ , by the above exact sequence (2.1) for  $\text{Cat} = \text{Top}$ , it suffices to prove  $q^*: [\mathbb{S}^{16}, \text{Top}/O] \rightarrow [X, \text{Top}/O]$  is monic, which is equivalent to saying that  $(S(g))^*: [S\mathbb{S}^8, \text{Top}/O] \rightarrow [\mathbb{S}^{16}, \text{Top}/O]$  is the zero homomorphism. For the case  $g = p$ , where  $p: \mathbb{S}^{15} \rightarrow \mathbb{S}^8$  is the Hopf map,  $(S(g))^*$  is the zero homomorphism, which was proved in the course of the proof of Lemma 1 in [2, pp.58–59]. This proof works verbatim for any map  $g: \mathbb{S}^{2n-1} \rightarrow \mathbb{S}^n$  as well. This proves (iii).

(iv): If  $n$  is any even integer such that  $\Theta_n$  and  $\Theta_{n+1}$  are trivial, then  $\pi_{n+1}(G/O) = 0$ . This shows that from the above exact sequence (2.1) for  $\text{Cat} = G$ ,  $q^*: [\mathbb{S}^{2n}, G/O] \rightarrow [X, G/O]$  is injective. Then  $f_M^* = \phi^* \circ q^*: [\mathbb{S}^{2n}, G/O] \rightarrow [M, G/O]$  is injective. From

the surgery exact sequences of  $M$  and  $\mathbb{S}^{2n}$ , we get the following commutative diagram ([3, Lemma 3.4]):

$$(2.2) \quad \begin{array}{ccccccc} L_{2n+1}(e) & \longrightarrow & \bar{\Theta}_{2n} & \xrightarrow{\eta_{\mathbb{S}^{2n}}} & \pi_{2n}(G/O) & \longrightarrow & L_{2n}(e) \\ \downarrow = & & \downarrow f_M^\bullet & & \downarrow f_M^* & & \downarrow = \\ L_{2n+1}(e) & \longrightarrow & \mathcal{S}^{Diff}(M) & \xrightarrow{\eta_M} & [M, G/O] & \longrightarrow & L_{2n}(e) \end{array}$$

By using the facts that  $L_{2n+1}(e) = 0$ , injectivity of  $\eta_{\mathbb{S}^{2n}}$  and  $\eta_M$  follow from the diagram, and combine with the injectivity of  $f_M^*$  to show that  $f_M^\bullet: \bar{\Theta}_{2n} \rightarrow \mathcal{S}^{Diff}(M)$  is injective.  $I_h(M)$  is exactly the kernel of  $f_M^\bullet$ , and so  $I_h(M) = 0$ . This proves (iv).  $\square$

REMARK 2.8. (i) By M. Kervaire and J. Milnor [6],  $\Theta_m = 0$  for  $m = 1, 2, 3, 4, 5, 6$  or  $12$ . If  $M^{2n}$  is a closed smooth  $(n-1)$ -connected  $2n$ -manifold, by Theorem 2.7 (i) and (ii),  $I_c(M^{2n}) = 0$  for  $n = 3, 4, 5$  or  $11$  and  $\mathcal{C}(M^{2n}) \cong \bar{\Theta}_{2n}$  for  $n = 4$  or  $5$ .  
 (ii) If  $M$  has the homotopy type of  $\mathbb{O}\mathbb{P}^2$ , by Theorem 1.1 and Theorem 2.7 (iii), we have  $I_c(M) = 0 \neq I(M)$ .  
 (iii) By Theorem 2.7 (iv), if  $M$  has the homotopy type of  $\mathbb{H}\mathbb{P}^2$ , then  $I_h(M) = 0$ .

DEFINITION 2.9. Let  $M$  and  $N$  are smooth manifolds. A smooth map  $f: M \rightarrow N$  is called tangential if for some integers  $k, l$ ,  $f^*(T(N)) \oplus \epsilon_M^k \cong T(M) \oplus \epsilon_M^l$ .

DEFINITION 2.10. Let  $M$  be a topological manifold. Let  $(N, f)$  be a pair consisting of a smooth manifold  $N$  together with a tangential homotopy equivalence of degree one  $f: N \rightarrow M$ . Two such pairs  $(N_1, f_1)$  and  $(N_2, f_2)$  are equivalent provided there exists a diffeomorphism  $g: N_1 \rightarrow N_2$  such that  $f_2 \circ g$  is homotopic to  $f_1$ . The set of all such equivalence classes is denoted by  $\theta(M)$ .

For  $M = \mathbb{H}\mathbb{P}^2$ , [5, Theorem 4] shows  $\theta(\mathbb{H}\mathbb{P}^2)$  contains at most two elements. Now by Remark 2.8 (iii), we have the following:

**Corollary 2.11.**  $\theta(\mathbb{H}\mathbb{P}^2)$  contains exactly two elements, with representatives given by  $(\mathbb{H}\mathbb{P}^2, \text{id})$  and  $(\mathbb{H}\mathbb{P}^2 \# \Sigma^8, \text{id})$ , where  $\Sigma^8$  is the exotic 8-sphere.

### 3. Inertia groups of projective plane-like manifolds

In [15], C.T.C. Wall assigned to each closed oriented  $(n-1)$ -connected  $2n$ -dimensional smooth manifold  $M^{2n}$  with  $n \geq 3$ , a system of invariants as follows:

- (1)  $H = H^n(M; \mathbb{Z}) \cong \text{Hom}(H_n(M; \mathbb{Z}), \mathbb{Z}) \cong \bigoplus_{j=1}^k \mathbb{Z}$ , the cohomology group of  $M$ , with  $k$  the  $n$ -th Betti number of  $M$ ,
- (2)  $I: H \times H \rightarrow \mathbb{Z}$ , the intersection form of  $M$  which is unimodular and  $n$ -symmetric, defined by

$$I(x, y) = \langle x \cup y, [M] \rangle,$$

where the homology class  $[M]$  is the orientation class of  $M$ ,

(3) A map  $\alpha: H^n(M; \mathbb{Z}) \rightarrow \pi_{n-1}(SO_n)$  that assigns each element  $x \in H^n(M; \mathbb{Z})$  to the characteristic map  $\alpha(x)$  for the normal bundle of the embedded  $n$ -sphere  $S_x^n$  representing  $x$ .

Denote by  $\chi = S \circ \alpha: H^n(M; \mathbb{Z}) \rightarrow \pi_{n-1}(SO_{n+1}) \cong \widetilde{KO}(S^n)$ , where  $S: \pi_{n-1}(SO_n) \rightarrow \pi_{n-1}(SO_{n+1})$  is the suspension map. Then

$$\chi = S \circ \alpha \in H^n(M; \widetilde{KO}(S^n)) = \text{Hom}(H^n(M; \mathbb{Z}); \widetilde{KO}(S^n))$$

can be viewed as an  $n$ -dimensional cohomology class of  $M$ , with coefficients in  $\widetilde{KO}(S^n)$ . The obstruction to triviality of the tangent bundle over the  $n$ -skeleton is the element  $\chi \in H^n(M; \widetilde{KO}(S^n))$  [15]. By [15, pp. 179–180], the Pontrjagin class of  $M^{2n}$  is given by

$$(3.1) \quad p_m(M^{2n}) = \pm a_m(2m - 1)! \chi,$$

where  $n = 4m$  and

$$a_m = \begin{cases} 1 & \text{if } 4m \equiv 0 \pmod{8}, \\ 2 & \text{if } 4m \equiv 4 \pmod{8}. \end{cases}$$

Define  $\Theta_n(k)$  to be the subgroup of  $\bar{\Theta}_n$  consisting of those homotopy  $n$ -sphere  $\Sigma^n$  which are the boundaries of  $k$ -connected  $(n + 1)$ -dimensional compact manifolds,  $1 \leq k < [n/2]$ . Thus,  $\Theta_n(k)$  is the kernel of the natural map  $i_k: \bar{\Theta}_n \rightarrow \Omega_n(k)$ , where  $\Omega_n(k)$  is the  $n$ -dimensional group in  $k$ -connective cobordism theory [13] and  $i_k$  sends  $\Sigma^n$  to its cobordism class. Using surgery, we see  $\Omega_{\ast}(1)$  is the usual oriented cobordism group. So  $\bar{\Theta}_n = \Theta_n(1)$ . Similarly,  $\Omega_n(2) \cong \Omega_n^{Spin}$  ( $n \geq 7$ ); since  $BSpin$  is, in fact, 3-connected, for  $n \geq 8$ ,  $\Omega_n(2) \cong \Omega_n(3)$  and  $\Theta_n(2) = \Theta_n(3) = bSpin_n$ . Here  $bSpin_n$  consists of homotopy  $n$ -sphere which bound spin manifolds.

In [16], C.T.C. Wall defined the Grothendieck group  $\mathcal{G}_n^{2n+1}$ , a homomorphism  $\vartheta: \mathcal{G}_n^{2n+1} \rightarrow \bar{\Theta}_{2n}$  such that  $\vartheta(\mathcal{G}_n^{2n+1}) = \Theta_{2n}(n - 1)$  and proved the following theorem:

**Theorem 3.1** (Wall). *Let  $M^{2n}$  be a closed smooth  $(n - 1)$ -connected  $2n$ -manifold and  $\Sigma^{2n}$  be a homotopy sphere in  $\bar{\Theta}_{2n}$ . Then  $M \# \Sigma^{2n}$  is an orientation-preserving diffeomorphic to  $M$  if and only if*

- (i)  $\Sigma^{2n} = 0$  in  $\bar{\Theta}_{2n}$  or
- (ii)  $\chi \not\equiv 0 \pmod{2}$  and  $\Sigma^{2n} \in \vartheta(\mathcal{G}_n^{2n+1}) = \Theta_{2n}(n - 1)$

We also need the following result from [1]:

**Theorem 3.2** (Anderson, Brown, Peterson). *Let  $\eta_n: \bar{\Theta}_n \rightarrow \Omega_n^{Spin}$  be the homomorphism such that  $\eta_n$  sends  $\Sigma^n$  to its spin cobordism class. Then  $\eta_n \neq 0$  if and only if  $n = 8k + 1$  or  $8k + 2$ .*

Proof of Theorem 1.1. Let  $\xi$  be a generator of  $H^n(M^{2n}; \mathbb{Z})$ . Consider the case  $n = 4$ . Then by Itiro Tamura [14] and (3.1), the Pontrjagin class of  $M^{2n}$  is given by

$$p_1(M^{2n}) = 2(2h + 1)\xi = \pm 2\chi,$$

where  $h \in \mathbb{Z}$ . This implies that

$$\chi = \pm(2h + 1)\xi.$$

Likewise, for  $n = 8$ , we have

$$p_2(M^{2n}) = 6(2k + 1)\xi = \pm 6\chi,$$

where  $k \in \mathbb{Z}$ . This implies that

$$\chi = \pm(2k + 1)\xi.$$

Therefore in either case,  $\chi \not\equiv 0 \pmod{2}$ . Now by Theorem 3.1, it follows that

$$I(M^{2n}) = \Theta_{2n}(n - 1).$$

Since  $\Theta_{2n}(n - 1)$  is the kernel of the natural map  $i_{n-1}: \bar{\Theta}_{2n} \rightarrow \Omega_{2n}(n - 1)$ , where  $\Omega_{2n}(n - 1) \cong \Omega_8^{Spin}$  for  $n = 4$  and  $\Omega_{2n}(n - 1) \cong \Omega_{16}^{String} \cong \mathbb{Z} \oplus \mathbb{Z}$  for  $n = 8$  [4]. Now by Theorem 3.2 and using the fact that  $\bar{\Theta}_{16} \cong \mathbb{Z}_2$  [6], we have  $i_{n-1} = 0$  for  $n = 4$  and  $8$ . This shows that  $\Theta_{2n}(n - 1) = \bar{\Theta}_{2n}$ . This implies that

$$I(M^{2n}) \cong \mathbb{Z}_2.$$

This completes the proof of Theorem 1.1. □

#### 4. Smooth structures of $(n - 1)$ -connected $2n$ -manifolds

DEFINITION 4.1 (*Cat = Diff* or *Top*-structure sets, [3]). Let  $M$  be a closed *Cat*-manifold. We define the *Cat*-structure set  $S^{Cat}(M)$  to be the set of equivalence classes of pairs  $(N, f)$  where  $N$  is a closed *Cat*-manifold and  $f: N \rightarrow M$  is a homotopy equivalence. And the equivalence relation is defined as follows:

$$(N_1, f_1) \sim (N_2, f_2) \text{ if there is a } Cat\text{-isomorphism } \phi: N_1 \rightarrow N_2 \text{ such that } f_2 \circ \phi \text{ is homotopic to } f_1.$$

We will denote the class in  $S^{Cat}(M)$  of  $(N, f)$  by  $[(N, f)]$ . The base point of  $S^{Cat}(M)$  is the equivalence class  $[(M, id)]$  of  $id: M \rightarrow M$ .

The forgetful maps  $F_{Diff}: S^{Diff}(M) \rightarrow S^{Top}(M)$  and  $F_{Con}: \mathcal{C}(M) \rightarrow S^{Diff}(M)$  fit into a short exact sequence of pointed sets [3]:

$$\mathcal{C}(M) \xrightarrow{F_{Con}} S^{Diff}(M) \xrightarrow{F_{Diff}} S^{Top}(M).$$

**Theorem 4.2.** *Let  $n$  be any integer greater than 3 such that  $\Theta_n$  and  $\Theta_{n+1}$  are trivial and  $M^{2n}$  be a closed smooth  $(n - 1)$ -connected  $2n$ -manifold. Let  $f: N \rightarrow M$  be a homeomorphism where  $N$  is a closed smooth manifold. Then*

(i) *there exists a diffeomorphism  $\phi: N \rightarrow M \# \Sigma^{2n}$ , where  $\Sigma^{2n} \in \bar{\Theta}_{2n}$  such that the following diagram commutes up to homotopy:*

$$\begin{array}{ccc} N & \xrightarrow{\phi} & M \# \Sigma^{2n} \\ & \searrow f & \downarrow \text{id} \\ & & M \end{array}$$

(ii) *If  $I_h(M) = \bar{\Theta}_{2n}$ , then  $f: N \rightarrow M$  is homotopic to a diffeomorphism.*

Proof. Consider the short exact sequence of pointed sets

$$\mathcal{C}(M) \xrightarrow{F_{Con}} \mathcal{S}^{Diff}(M) \xrightarrow{F_{Diff}} \mathcal{S}^{Top}(M).$$

By Theorem 2.7 (ii), we have

$$\mathcal{C}(M) = \{[M \# \Sigma] \mid \Sigma \in \bar{\Theta}_{2n}\} \cong \bar{\Theta}_{2n}.$$

Since  $[(N, f)] \in F_{Diff}^{-1}([(M, \text{id})])$ , we obtain

$$[(N, f)] \in \text{Im}(F_{Con}) = \{[M \# \Sigma] \mid \Sigma \in \bar{\Theta}_{2n}\}.$$

This implies that there exists a homotopy sphere  $\Sigma^{2n} \in \bar{\Theta}_{2n}$  such that  $(N, f) \sim (M \# \Sigma^{2n}, \text{id})$  in  $\mathcal{S}^{Diff}(M)$ . This implies that there exists a diffeomorphism  $\phi: N \rightarrow M \# \Sigma^{2n}$  such that  $f$  is homotopic to  $\text{id} \circ \phi$ . This proves (i).

If  $I_h(M) = \bar{\Theta}_{2n}$ , then  $\text{Im}(F_{Con}) = \{[(M, \text{id})]\}$  and hence  $(N, f) \sim (M, \text{id})$  in  $\mathcal{S}^{Diff}(M)$ . This shows that  $f: N \rightarrow M$  is homotopic to a diffeomorphism  $N \rightarrow M$ . This proves (ii). □

**Theorem 4.3.** *Let  $n$  be any integer greater than 3 such that  $\Theta_n$  and  $\Theta_{n+1}$  are trivial and  $M^{2n}$  be a closed smooth  $(n - 1)$ -connected  $2n$ -manifold. Then the number of distinct smooth structures on  $M^{2n}$  up to diffeomorphism is less than or equal to the cardinality of  $\bar{\Theta}_{2n}$ . In particular, the set of diffeomorphism classes of smooth structures on  $M^{2n}$  is  $\{[M \# \Sigma] \mid \Sigma \in \bar{\Theta}_{2n}\}$ .*

Proof. By Theorem 4.2 (i), if  $N$  is a closed smooth manifold homeomorphic to  $M$ , then  $N$  is diffeomorphic to  $M \# \Sigma^{2n}$  for some homotopy  $2n$ -sphere  $\Sigma^{2n}$ . This implies that the set of diffeomorphism classes of smooth structures on  $M^{2n}$  is  $\{[M \# \Sigma] \mid \Sigma \in \bar{\Theta}_{2n}\}$ . This shows that the number of distinct smooth structures on  $M^{2n}$  up to diffeomorphism is less than or equal to the cardinality of  $\bar{\Theta}_{2n}$ . □



REMARK 4.4. (1) By Theorem 4.3, every closed smooth 3-connected 8-manifold has at most two distinct smooth structures up to diffeomorphism.

(2) If  $M^8$  is a closed smooth 3-connected 8-manifold such that  $H^4(M; \mathbb{Z}) \cong \mathbb{Z}$ , then by Theorem 1.1,  $I(M) \cong \mathbb{Z}_2$ . Now by Theorem 4.3,  $M$  has a unique smooth structure up to diffeomorphism.

(3) If  $M = \mathbb{S}^4 \times \mathbb{S}^4$ , then by Theorem 4.3,  $\mathbb{S}^4 \times \mathbb{S}^4$  has at most two distinct smooth structures up to diffeomorphism, namely,  $\{[\mathbb{S}^4 \times \mathbb{S}^4], [\mathbb{S}^4 \times \mathbb{S}^4 \# \Sigma]\}$ , where  $\Sigma$  is the exotic 8-sphere. However, by [11, Theorem A],  $I(\mathbb{S}^4 \times \mathbb{S}^4) = 0$ . This implies that  $\mathbb{S}^4 \times \mathbb{S}^4$  has exactly two distinct smooth structures.

**Theorem 4.5.** *Let  $M$  be a closed smooth 3-connected 8-manifold with stable tangential invariant  $\chi = S \circ \alpha: H_4(M; \mathbb{Z}) \rightarrow \pi_3(SO) = \mathbb{Z}$ . Then  $M$  has exactly two distinct smooth structures up to diffeomorphism if and only if  $\text{Im}(S \circ \alpha) \subseteq 2\mathbb{Z}$ .*

Proof. Suppose  $M$  has exactly two distinct smooth structures up to diffeomorphism. Then by Theorem 4.3,  $M$  and  $M \# \Sigma$  are not diffeomorphic, where  $\Sigma$  is the exotic 8-sphere. Since  $\bar{\Theta}_8 = \Theta_8(3)$ , by Theorem 3.1, the stable tangential invariant  $\chi$  is zero (mod 2) and hence  $\text{Im}(S \circ \alpha) \subseteq 2\mathbb{Z}$ . Conversely, suppose  $\text{Im}(S \circ \alpha) \subseteq 2\mathbb{Z}$ . Now by Theorem 3.1,  $M$  can not be diffeomorphic to  $M \# \Sigma$ , where  $\Sigma$  is the exotic 8-sphere. Now by Theorem 4.3,  $M$  has exactly two distinct smooth structures up to diffeomorphism.  $\square$

REMARK 4.6. If  $n = 2, 3, 5, 6, 7 \pmod{8}$  or the stable tangential invariant  $\chi$  of  $M^{2n}$  is zero (mod 2), then by [16, Corollary, p.289] and Theorem 3.1, we have  $I(M^{2n}) = 0$ . So, by Theorem 4.3, we have the following:

**Theorem 4.7.** *Let  $n$  be any integer greater than 3 such that  $\Theta_n$  and  $\Theta_{n+1}$  are trivial and  $M^{2n}$  be a closed smooth  $(n - 1)$ -connected  $2n$ -manifold. If  $n = 2, 3, 5, 6, 7 \pmod{8}$  or the stable tangential invariant  $\chi$  of  $M^{2n}$  is zero (mod 2), then the set of diffeomorphism classes of smooth structures on  $M^{2n}$  is in one-to-one correspondence with group  $\bar{\Theta}_{2n}$ .*

REMARK 4.8. (1) By Theorem 4.7, every closed smooth 4-connected 10-manifold has exactly six distinct smooth structures, namely,  $\{[M \# \Sigma] \mid \Sigma \in \bar{\Theta}_{10} \cong \mathbb{Z}_6\}$ .

(2) If  $M^{2n}$  is  $n$ -parallelisable, almost parallelisable or  $\pi$ -manifold, then the stable tangential invariant  $\chi$  of  $M$  is zero [15]. Then by Theorem 4.7, we have the following:

**Corollary 4.9.** *Let  $n$  be any integer greater than 3 such that  $\Theta_n$  and  $\Theta_{n+1}$  are trivial and  $M^{2n}$  be a closed smooth  $(n - 1)$ -connected  $2n$ -manifold. If  $M^{2n}$  is  $n$ -parallelisable, almost parallelisable or  $\pi$ -manifold, then the set of diffeomorphism classes of smooth structures on  $M^{2n}$  is in one-to-one correspondence with group  $\bar{\Theta}_{2n}$ .*

DEFINITION 4.10 ([8]). The normal  $k$ -type of a closed smooth manifold  $M$  is the fibre homotopy type of a fibration  $p: B \rightarrow BO$  such that the fibre of the map  $p$

is connected and its homotopy groups vanish in dimension  $\geq k + 1$ , admitting a lift of the normal Gauss map  $\nu_M: M \rightarrow BO$  to a map  $\bar{\nu}_M: M \rightarrow B$  such that  $\bar{\nu}_M: M \rightarrow B$  is a  $(k + 1)$ -equivalence, i.e., the induced homomorphism  $\bar{\nu}_M: \pi_i(M) \rightarrow \pi_i(B)$  is an isomorphism for  $i \leq k$  and surjective for  $i = k + 1$ . We call such a lift a normal  $k$ -smoothing.

**Theorem 4.11.** *Let  $n = 5, 7$  and let  $M_0$  and  $M_1$  be closed smooth  $(n - 1)$ -connected  $2n$ -manifolds with the same Euler characteristic. Then*

- (i) *There is a homotopy sphere  $\Sigma^{2n} \in \bar{\Theta}_{2n}$  such that  $M_0$  and  $M_1 \# \Sigma^{2n}$  are diffeomorphic.*
- (ii) *Let  $M^{2n}$  be a closed smooth  $(n - 1)$ -connected  $2n$ -manifold such that  $[M] = 0 \in \Omega_{2n}^{String}$  and let  $\Sigma$  be any exotic  $2n$ -sphere in  $\bar{\Theta}_{2n}$ . Then  $M$  and  $M \# \Sigma$  are not diffeomorphic.*

Proof. (i):  $M_0$  and  $M_1$  are  $(n - 1)$ -connected, and  $n$  is 5 or 7; therefore,  $p_1/2$  and the Stiefel–Whitney classes  $\omega_2$  vanish. So,  $M_0$  and  $M_1$  are  $BString$ -manifolds. Let  $\bar{\nu}_{M_j}: M_j \rightarrow BString$  be a lift of the normal Gauss map  $\nu_{M_j}: M_j \rightarrow BO$  in the fibration  $p: BString = BO\langle 8 \rangle \rightarrow BO$ , where  $j = 0$  and 1. Since  $BString$  is 7-connected,  $p_\#: \pi_i(BString) \rightarrow \pi_i(BO)$  is an isomorphism for all  $i \geq 8$ . This shows that  $\bar{\nu}_{M_j}: M_j \rightarrow BString$  is an  $n$ -equivalence and hence the normal  $(n - 1)$ -type of  $M_0$  and  $M_1$  is  $p: BString \rightarrow BO$ . We know that  $\Omega_{2n}^{String} \cong \bar{\Theta}_{2n}$ , where the group structure is given by connected sum [4]. This implies that there always exists  $\Sigma^{2n} \in \bar{\Theta}_{2n}$  such that  $M_0$  and  $M_1 \# \Sigma^{2n}$  are  $BString$ -bordant. Since  $M_0$  and  $M_1 \# \Sigma^{2n}$  have the same Euler characteristic, by [8, Corollary 4],  $M_0$  and  $M_1 \# \Sigma^{2n}$  are diffeomorphic.

(ii): Since the image of the standard sphere under the isomorphism  $\bar{\Theta}_{2n} \cong \Omega_{2n}^{String}$  represents the trivial element in  $\Omega_{2n}^{String}$ , we have  $[M^{2n}] \neq [M \# \Sigma]$  in  $\Omega_{2n}^{String}$ . This implies that  $M$  and  $M \# \Sigma$  are not  $BString$ -bordant. By obstruction theory,  $M^{2n}$  has a unique string structure. This implies that  $M$  and  $M \# \Sigma$  are not diffeomorphic.  $\square$

**Theorem 4.12.** *Let  $M$  be a closed smooth 6-connected 14-dimensional  $\pi$ -manifold and  $\Sigma$  is the exotic 14-sphere. Then  $M \# \Sigma$  is not diffeomorphic to  $M$ . Thus,  $I(M) = 0$ . Moreover, if  $N$  is a closed smooth manifold homeomorphic to  $M$ , then  $N$  is diffeomorphic to either  $M$  or  $M \# \Sigma$ .*

Proof. It follows from results of Anderson, Brown and Peterson on spin cobordism [1] that the image of the natural homomorphism  $\Omega_{14}^{framed} \rightarrow \Omega_{14}^{Spin}$  is 0 and  $\Omega_{14}^{String} \cong \Omega_{14}^{Spin} \cong \mathbb{Z}_2$  [4]. This shows that  $[M] = 0 \in \Omega_{14}^{String}$ . Now by Theorem 4.11 (ii),  $M \# \Sigma$  is not diffeomorphic to  $M$ . If  $N$  is a closed smooth manifold homeomorphic to  $M$ , then  $N$  and  $M$  have the same Euler characteristic. Then by Theorem 4.11 (i),  $N$  is diffeomorphic to either  $M$  or  $M \# \Sigma$ .  $\square$

REMARK 4.13. By the above Theorem 4.12, the set of diffeomorphism classes of smooth structures on a closed smooth 6-connected 14-dimensional  $\pi$ -manifold  $M$  is

$$\{[M], [M \# \Sigma]\} \cong \mathbb{Z}_2,$$

where  $\Sigma$  is the exotic 14-sphere. So, the number of distinct smooth structures on  $M$  is 2.

ACKNOWLEDGMENTS. The author would like to thank his advisor, Prof. A.R. Shastri for several helpful suggestions and questions.

---

### References

- [1] D.W. Anderson, E.H. Brown, Jr. and F.P. Peterson: *The structure of the Spin cobordism ring*, Ann. of Math. (2) **86** (1967), 271–298.
- [2] C.S. Aravinda and F.T. Farrell: *Exotic negatively curved structures on Cayley hyperbolic manifolds*, J. Differential Geom. **63** (2003), 41–62.
- [3] D. Crowley: *The smooth structure set of  $S^p \times S^q$* , Geom. Dedicata **148** (2010), 15–33.
- [4] V. Giambalvo: *On  $\{8\}$ -cobordism*, Illinois J. Math. **15** (1971), 533–541.
- [5] D.N. Hertz: *Ambient surgery and tangential homotopy quaternionic projective spaces*, Trans. Amer. Math. Soc. **145** (1969), 517–545.
- [6] M.A. Kervaire and J.W. Milnor: *Groups of homotopy spheres, I*, Ann. of Math. (2) **77** (1963), 504–537.
- [7] A. Kosinski: *On the inertia group of  $\pi$ -manifolds*, Amer. J. Math. **89** (1967), 227–248.
- [8] M. Kreck: *Surgery and duality*, Ann. of Math. (2) **149** (1999), 707–754.
- [9] L. Kramer and S. Stolz: *A diffeomorphism classification of manifolds which are like projective planes*, J. Differential Geom. **77** (2007), 177–188.
- [10] R.C. Kirby and L.C. Siebenmann: *Foundational Essays on Topological Manifolds, Smoothings, and Triangulations*, Ann. of Math. Studies **88**, Princeton Univ. Press, Princeton, NJ, 1977.
- [11] R. Schultz: *On the inertia group of a product of spheres*, Trans. Amer. Math. Soc. **156** (1971), 137–153.
- [12] S. Smale: *On the structure of manifolds*, Amer. J. Math. **84** (1962), 387–399.
- [13] R.E. Stong: *Notes on Cobordism Theory*, Princeton Univ. Press, Princeton, NJ, 1968.
- [14] I. Tamura: *8-manifolds admitting no differentiable structure*, J. Math. Soc. Japan **13** (1961), 377–382.
- [15] C.T.C. Wall: *Classification of  $(n - 1)$ -connected  $2n$ -manifolds*, Ann. of Math. (2) **75** (1962), 163–189.
- [16] C.T.C. Wall: *Classification problems in differential topology. VI. Classification of  $(s - 1)$ -connected  $(2s + 1)$ -manifolds*, Topology **6** (1967), 273–296.

Department of Mathematics  
 Indian Institute of Technology, Bombay  
 Mumbai-76  
 India  
 e-mail: rameshk@math.iitb.ac.in  
 e-mail: mathsramesh1984@gmail.com